

<https://helda.helsinki.fi>

---

## High prevalence of Clostridium botulinum in vegetarian sausages

Pernu, Noora

2020-10

---

Pernu , N , Keto-Timonen , R , Lindström , M & Korkeala , H 2020 , ' High prevalence of Clostridium botulinum in vegetarian sausages ' , Food Microbiology , vol. 91 , 103512 . <https://doi.org/10.1016/j.fm.2020.103512>

---

<http://hdl.handle.net/10138/317886>

<https://doi.org/10.1016/j.fm.2020.103512>

---

cc\_by

acceptedVersion

---

*Downloaded from Helda, University of Helsinki institutional repository.*

*This is an electronic reprint of the original article.*

*This reprint may differ from the original in pagination and typographic detail.*

*Please cite the original version.*

# Journal Pre-proof

High prevalence of *Clostridium botulinum* in vegetarian sausages

Noora Pernu, Riikka Keto-Timonen, Miia Lindström, Hannu Korkeala

PII: S0740-0020(20)30101-5

DOI: <https://doi.org/10.1016/j.fm.2020.103512>


Reference: YFMIC 103512

To appear in: *Food Microbiology*

Received Date: 21 December 2019

Revised Date: 4 April 2020

Accepted Date: 12 April 2020



Please cite this article as: Pernu, N., Keto-Timonen, R., Lindström, M., Korkeala, H., High prevalence of *Clostridium botulinum* in vegetarian sausages, *Food Microbiology* (2020), doi: <https://doi.org/10.1016/j.fm.2020.103512>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier Ltd.

Short communication

Title: High prevalence of *Clostridium botulinum* in vegetarian sausages

Noora Pernu<sup>1</sup>, Riikka Keto-Timonen<sup>1</sup>, Miia Lindström\*, and Hannu Korkeala

Department of Food Hygiene and Environmental Health, Faculty of Veterinary Medicine, P.O. Box 66, FI-00014 University of Helsinki, Finland

<sup>1</sup>These authors contributed equally to this work.\*Author for correspondence: Miia Lindström, tel. +358504486587, fax +358294157101, e-mail [miia.lindstrom@helsinki.fi](mailto:miia.lindstrom@helsinki.fi)

## 1. Introduction

In line with sustainable development goals, plant-based foods and meat substitutes are becoming a preferred source of protein among increasing number of consumers. A popular alternative for meat products are vegetarian sausages, with a range of products being available on the market. Most products are vacuum-packaged chilled foods, but also frozen and canned vegetarian sausages are retailed. Ingredients include typically a plant or fungal protein source (soy, wheat protein, chickpea, pea protein, mycoprotein), other vegetables (corn, potato, pepper, tomato, onion, garlic etc.), herbs, spices, salt, vegetable oil, and additives (thickening agents, stabilizers, pH regulators, antioxidants).

A major food safety concern in vacuum-packaged chilled foods are psychrotrophic, botulinum neurotoxin (BoNT) producing clostridia, particularly *Clostridium botulinum* Group II (Lindström et al., 2006a; Peck, 2006). These bacteria produce resistant endospores, grow in anaerobic conditions, and produce the highly potent BoNTs during growth. Once ingested, BoNTs inhibit the release of acetylcholine at the neuromuscular junction and cause a potentially lethal flaccid paralysis, botulism. *C. botulinum* spores exist widely in environment and can contaminate food raw materials. The spores can survive pasteurization and, under favorable conditions, may germinate and outgrow into toxic cultures.

Both home-canned and commercially processed vegetables are common sources of foodborne botulism (Sobel et al., 2004; Anniballi et al., 2017; Hellmich et al., 2018). A range of vegetables have been implicated in outbreaks, including onions, potatoes, corn, peppers, asparagus, carrots, beans, olives, and garlic (MacDonald et al., 1985; Morse et al., 1990; Angulo et al., 1998; Sobel et al., 2004; Zanon et al., 2006; Date et al., 2011; Jalava et al., 2011; Lindström et al., 2011; Hill et al., 2013). In addition, tofu has caused botulism, introducing soy into the list of implicated vegetables (Chai et al., 2013). However, screening studies on the prevalence of *C. botulinum* spores in non-outbreak-related vegetables are scarce. Negligible positive findings (sample size ranging between 10-200 g) suggest a very low prevalence and spore contamination levels below 1–10 spores/kg (Insalata et al., 1969; Lilly et al., 1995; Braconnier et al., 2001;

Sevenier et al., 2012; Barker et al., 2016). The few positive screening samples and most outbreak investigations revealed *C. botulinum* types A and B. While type A strains are exclusively of the mesophilic Group I, which do not possess a risk of growth under refrigeration, BoNT B-producing strains may belong to either Group I or Group II and can be distinguished by metabolic features (Dahlsten et al., 2008) and by the neurotoxin gene sequence (Hill et al., 2007). Unfortunately, information on the physiological group or toxin gene sequence for early *C. botulinum* type B findings in vegetables are not available, but a recent report on a subtype B4 neurotoxin gene present in two out of three studied samples of porcini mushrooms (Barker et al., 2016) confirms that the psychrotrophic Group II strains may be of concern. This is in line with our previous finding of Group II *C. botulinum* type E in a sample of vegetarian sausage (Lindström et al., 2001).

Vegetarian sausages can be categorized as refrigerated processed foods of extended durability (REPFED). Typically, REPFEDs are processed at mild pasteurization temperatures, cooled rapidly after processing, and stored refrigerated over extended periods of time (Gorris and Peck, 1998). The safety of REPFEDs relies on hurdle technology combining multiple preservation factors to control microbial growth (Gorris and Peck, 1998). The applied heat treatments, prevailing storage temperatures, and the use of preservatives define the product shelf-life and safety.

While there are no reported cases of botulism due to vegetarian sausages, the possibility of raw material contamination with *C. botulinum* Groups I and II spores, mild pasteurization, vacuum-packaging, and long shelf-lives contribute to an apparently high risk of *C. botulinum* growth and BoNT production related to vacuum-packaged chilled vegetarian sausages. Here we show a high overall prevalence of 32% of *C. botulinum* in 74 vegetarian sausage products.

## 2. Material and methods

### 2.1 Vegetarian sausages.

A total of 74 samples of frozen (8) or chilled (66) packaged vegetarian sausages from seven producers were purchased in Finland and Germany. The pH of such products is above 5.7 and added NaCl concentrations in the range of 1.2–1.9%. Assuming dry-matter concentrations of 27–68% (Havlik et al., 2010), the corresponding water-phase NaCl concentrations are mainly in the range of 2–4%, and exceed 5% only in the rare occasions when added NaCl concentration exceeds 1.9% and dry-matter content exceeds 64%. The shelf-lives of the investigated vacuum-packaged products remaining at the time of purchase varied from less than 2 weeks to 6 months. Some of the investigated products contained detailed instructions for cooking, including heating temperature and time, some products contained just a suggestion of heating method without time indications, and some products were advised to be served either heated or cold. The main ingredients of the vegetarian sausages were soy (soy protein or tofu), wheat protein, vegetable oil, sugar, spices, salt, and corn, wheat, or potato starch. Additives such as pH regulators, emulsifiers, stabilizers, and antioxidants were commonly included but were not identified in more detail. Some of the products contained also oat, rice, egg white, apple, onion and/or garlic.

## 2.2 Microbial analyses.

Before laboratory analysis, the vegetarian sausage samples were stored at temperatures instructed by the manufacturers, either frozen or at refrigeration. The quantity of *C. botulinum* was determined from non-heat-shocked samples using the most probable number (MPN) method (Cochran, 1950), using PCR detection of *C. botulinum* growth and the formula of Thomas (1942) for MPN estimation based on the number of PCR-positive tubes (Hielm et al., 1996). A sample size of 20 to 111 g was inoculated into a set of tubes containing tryptone–peptone–glucose–yeast extract (TPGY) broth (1:10) and incubated under anaerobic conditions at 30°C (Group II) or at 37°C (Group I) for 72–96 hours, followed by overnight cultures in fresh TPGY (1:10) under identical conditions. Cells from 1-ml aliquots were prepared for PCR templates as described (Lindström et al., 2001). The presence of genes encoding BoNT types A, B, E, and F was determined using multiplex PCR (Lindström et al., 2001). Attempts to isolate *C. botulinum* from all PCR-positive samples were made on egg yolk agar plates (Hauschild and Hilsheimer, 1977), and amplified

fragment length polymorphism (AFLP) method was used to genotype the *C. botulinum* isolates (Keto-Timonen *et al.*, 2006).

### 3. Results

We show a strikingly high overall prevalence of 32% for *C. botulinum* in vacuum-packaged vegetarian sausages (Table 1). Apart from one positive frozen sample (13%), all other positive samples were detected among chilled products (23 samples, 35%) with advised maximum storage temperatures of 6–10°C. Genes for BoNT types A, B, E, and F were detected, with types B (33 %), A (33%), and E (25%) detected frequently, and types B and F together, and types B and E together, once (4%, Table 1). The highest MPN counts were detected for Group II type E, up to 1200 cells/kg. At the time of purchase, eight samples had remaining shelf-lives of 3 to 6 months, and six of them were PCR-positive for *C. botulinum*.

Eight isolates were recovered from eight PCR-positive samples (MPN counts in these samples were 30-110 cells/kg): two type A, five type B, and one type E isolate. AFLP typing showed all the A and B isolates to be of *C. botulinum* Group I and the one E isolate to be of Group II (Fig. 1.). Successful isolation of *C. botulinum* validated the positive PCR findings. With successful recovery of a BoNT gene-carrying isolate as the sole criterion of a positive sample, the overall prevalence of *C. botulinum* in the vegetarian sausages appeared to be 11%.

### 4. Discussion

The MPN counts of *C. botulinum* in the vegetarian sausage samples varied between 20 to 1200 cells or spores/kg (average/median cell or spore count of 176/110 MPN/kg). Such counts are up to 3 log-units higher than the predicted counts of 1–10 spores/kg in raw materials (Barker *et al.*, 2016), and one log-unit higher than counts found in vacuum-packaged hot-smoked fish products (Hyytiä *et al.*, 1998) relatively frequently linked with botulism outbreaks (Kautter, 1964; Korkeala *et al.*, 1998; Lindström *et al.*, 2006b; King *et al.*, 2009). We found the highest *C. botulinum* counts ( $10^3$  cells or spores/kg) for *C. botulinum* Group

II type E, which are psychrotrophic and can grow and produce BoNT at temperatures as low as 3°C within 8 weeks (Graham et al., 1997). The high counts were detected in products with remaining shelf-lives at 6–10°C of up to 6 months. Thus, the safety risk related to these vacuum-packaged vegetarian sausages appears to be high.

The origin of *C. botulinum* in vegetarian sausages remains partially unclear. While vegetables are a common source for Group I *C. botulinum* types A and B, Group II type B strains are associated with pork meat preparations and occasionally to fish and seafood (Galazka and Przybylska, 1999; Lindström et al., 2006a; Mazuet et al., 2018). Type E is mostly associated with fish and seafood (Lindström et al., 2006a). It remains to be discussed how *C. botulinum* types less frequently associated with vegetables contaminate vegetarian sausages. A possible source for type E could be sea-salt as *C. botulinum* type E strains prevail in aquatic ecosystems (Hielm et al. 1998) and their spores have been detected in sea-salt (Fenicia et al., 2002).

While BoNT/A is exclusively produced by *C. botulinum* Group I strains and BoNT/E by Group II, BoNT/B and BoNT/F toxins can be produced by both *C. botulinum* Group I and II strains. The applied PCR methodology does not distinguish between BoNT/B or BoNT/F genes from Group I and II strains, and Group identification was obtained only when isolation of *C. botulinum* was successful. Indeed, AFLP typing showed all the five *bont/B*-positive isolates to belong to *C. botulinum* Group I. It remains unclear if the *bont/B*-positive and *bont/F*-positive samples that did not yield *C. botulinum* isolates were due to Group I or Group II. However, the similar prevalence and counts of *bont/A* and *bont/E* findings demonstrate that both Group I and Group II strains are prevalent in these products, and the applied heat treatments did not eliminate any of the two Groups. How the vegetarian sausages support the growth of Group I and II *C. botulinum* needs to be established.

Since the applied PCR assay (Lindström et al., 2001) fails to detect the genes for BoNT subtypes A2, A3 and A4 (De Medici et al., 2009), we may have underestimated the prevalence of *C. botulinum* type A. The only



type F-positive sample was also positive for type B, either due to the presence of two distinct strains each representing one of the two toxinotypes, or caused by a single bivalent type BF strain (Gimenez and Gimenez, 1993; Barash and Arnon, 2004). Another dual-positive sample yielded PCR signals for toxin types B and E. No bivalent type BE strains have been described in literature, thus this result is likely caused by the presence of two strains. Unfortunately, these two samples did not yield positive isolates for further analysis.

While *C. botulinum* Group I strains are of food safety concern due to the high heat resistance of their spores, their growth and toxin production can be controlled by refrigeration. However, Group II strains, whose spores are of moderate heat resistance, are of concern due to their growth and toxinogenic potential at low temperature (Graham et al., 1997; Derman et al., 2011). The time to toxinogenesis varies greatly by multiple factors, including the number of spores present, efficiency of the heat treatment applied, storage temperature, and intrinsic factors like water activity and pH (Segner et al., 1966; Hauschild and Hilsheimer, 1979; Meng and Genigeorgis, 1993; Peck et al., 1995). In previous inoculation studies, puréed mushrooms supported visible growth and toxin production in 20 days at 5°C (Carlin and Peck, 1996). At 3°C, growth and toxinogenesis were observed in a laboratory medium within 35 days (Graham et al., 1997). In the scenario of product shelf-life of 6 months (180 days) or longer, maintaining the storage temperature of vegetarian sausages strictly below 2.5°C (Graham et al., 1997) throughout the entire product lifespan appears imperative. Inoculation studies are needed to establish the growth potential from *C. botulinum* Group II spores in vegetarian sausages during storage at the temperatures of 6–8°C commonly applied for chilled foods. There is no information available on the growth and toxinogenic potential of *C. botulinum* Group II at 0–3°C for extended time periods of 6–9 months. This information appears pivotal for estimating the safety of extremely long shelf-lives for chilled products with obvious risk of *C. botulinum* growth.

To control the safety of vacuum-packaged chilled foods with shelf-lives over 10 days, it is advised that the products are given a 6D heat treatment, a process that will reduce the risk of *C. botulinum* Group II spores by a factor of  $10^6$  (ACMSF, 1992; ECFF, 2006). A temperature-time combination of 90°C 10 min has been proposed to provide a 6-log reduction (ACMSF, 1992; ECFF, 2006). Detection of  $10^2$ – $10^3$  *C. botulinum* Group II (type E) cells or spores/kg in the vacuum-packaged vegetarian sausages suggests that the pasteurization processes used to prepare these products are substantially less efficient than a 6D kill. Moreover, matrices containing lysozyme or other lytic enzymes might assist sublethally injured spores to germinate (Peck et al., 1993), thus heat treatments exceeding 90°C 10 min might be required to achieve a 6D kill (Peck and Fernandez, 1995). Vegetarian sausages may contain hen egg white, which is a rich source of lysozyme (Lund and Peck, 1994). Also plant-based lysozyme activity has been measured in wheat and maize (Lund and Peck, 1994). Whether these lytic activities challenge the heat treatments used in the production of vegetarian sausages, needs to be established. It is recommended that additional factors like chilled storage below 3°C, water activity below 0.97, corresponding to water-phase NaCl concentration of 5% or higher, or pH below 5.0 be applied to control spore germination and outgrowth (ACMSF, 1992). Since the cold chain cannot be controlled in the consumer domain, and the product pH is usually above 5.0 and the water-phase NaCl concentrations are mainly below 4%, the use of preservatives appears as an important means to control product safety.

Nitrite is a well-established antibotulinal agent used to cure meats (Kim and Foegeding, 1992, Keto-Timonen et al., 2012). However, its use is restricted in most non-meat products. While some leafy vegetables like lettuce and spinach and some root vegetables can contain significant amounts of nitrates that are reduced to nitrite during storage, soybean and soy products contain only 0.1 mg/kg nitrite (Kalaycioğlu and Erim, 2019). Thus natural nitrite levels in vegetarian sausages are supposed to be low (Kalaycioğlu and Erim, 2019).

Salts of organic acids have potential in controlling *C. botulinum* Group I and/or II growth. Potassium sorbate, sodium lactate, and sodium acetate at concentrations of 2–6% have growth-inhibitory potential (Seward et al., 1982; Kim and Foegeding, 1992; Miller et al., 1993; Meng and Genigeorgis, 1994) and are used in some vegetarian sausage products on the market. With the maximum concentration of 1000 mg/kg permitted in foods, sorbic acid also controlled *C. botulinum* Group I growth at pH below 5.5 (Lund et al. 1987). The effects of preservatives targeted to tackle with the product pH need to be properly established for soy-based products: in the presence of soy, *C. botulinum* Group I has been shown to grow and produce toxin at a pH as low as 4.1 (Young-Perkins and Merson, 1987) as opposed to the generally referred growth-inhibitory pH of 4.6 (Hauschild et al., 1975; Peck, 2006).

If measures fail to control *C. botulinum* growth and toxin production, BoNT can be destroyed by heating at 85°C for 5 minutes or at 80°C for 20 minutes (Woodburn et al., 1979). Vegetarian sausages are likely heated prior to consumption; however, not all products contain instructions for cooking. Moreover, some products are advised for consumption as cold, and light heating regimes might not destroy all preformed BoNT. Thus the safety of vegetarian sausage products must rely on multiple hurdles controlling growth and toxin production, and never merely on toxin inactivation during cooking.

In conclusion, a high prevalence of *C. botulinum* in vegetarian sausages suggests that these products could be a potential source of botulism. Mild heat treatments enable survival of *C. botulinum* Group I and Group II spores, and long shelf-lives may support spore germination, outgrowth and toxinogenesis. Inoculated pack studies and shelf-life tests are required to determine the growth and toxic potential of *C. botulinum* in vegetarian sausages and the length of safe shelf-lives.

226

227

228 **Acknowledgements**

229 This work was financially supported by the University of Helsinki and by the food industry. We thank Hanna  
230 Korpunen for her excellent technical assistance.

231

232 **References**

233 ACMSF (Advisory Committee on the Microbial Safety of Foods), 1992. Report on vacuum packing and  
234 associated process. Her Majesty's Stationery Office, London.

235

236 Angulo, F.J., Getz, J., Taylor, J.P., Hendricks, K.A., Hatheway, C.L., Barth, S.S., Solomon, H.M., Larson, A.E.,  
237 Johnson, E.A., Nickey, L.N., Ries, A.A., 1998. A large outbreak of botulism: the hazardous baked potato. J.  
238 Infect. Dis. 178(1), 172–177.

239

240 Anniballi, F., Auricchio, B., Fiore, A., Lonati, D., Locatelli, C.A., Lista, F., Fillo, S., Mandarino, G., De Medici, D.,  
241 2017. Botulism in Italy, 1986 to 2015. Euro Surveill. 22(24), 30550.

242

243 Barash, J.R., Arnon, S.S., 2004. Dual toxin-producing strain of *Clostridium botulinum* type Bf isolated from a  
244 California patient with infant botulism. J. Clin. Microbiol. 42(4), 1713–1715.

245

246 Barker, G.C., Malakar, P.K., Plowman, J., Peck, M.W., 2016. Quantification of nonproteolytic  
247 *Clostridium botulinum* spore loads in food materials. Appl. Environ. Microbiol. 82(6), 1675–1685.

248

249 Braconnier, A., Broussolle, V., Perelle, S., Fach, P., Nguyen-The, C., Carlin, F., 2001. Screening for  
250 *Clostridium botulinum* type A, B, and E in cooked chilled foods containing vegetables and raw material using  
251 polymerase chain reaction and molecular probes. J. Food Prot. 64(2), 201–207.

252

253 Carlin, F., Peck, M.W., 1996. Growth of and toxin production by nonproteolytic *Clostridium botulinum* in  
 254 cooked puréed vegetables at refrigeration temperatures. Appl Environ Microbiol. 62(8), 3069–3072.

255

256 Chai, E., Choi, E., Guitierrez, C., Melvin Hochman, M., Johnkutty, S., Kamel, W., Mekles, T., Zarnegar, R.,  
 257 Ackelsberg, J., Balter, S., Lee, E.H., Li, L., Ramos, A., Rodriguez, T., Weiss, D., Yung, J., Zhao, B., Davis, S.W.,  
 258 Egan, Hannett, G.E., Rao, A., Toprani, A., Sreenivasan, N., 2013. Botulism associated with home-fermented  
 259 tofu in two Chinese immigrants — New York City, March–April 2012. MMWR Morb Mortal Wkly Rep.  
 260 62(26), 529–532.

261

262 Cochran, W., 1950. Estimation of bacterial densities by means of the "Most Probable Number". Biometrics,  
 263 6(2), 105–116.

264

265 Dahlsten, E., Korkeala, H., Somervuo, P., Lindström, M., 2008. PCR assay for differentiating between Group I  
 266 (proteolytic) and Group II (nonproteolytic) strains of *Clostridium botulinum*. Int. J. Food Microbiol. 124(1),  
 267 108–111.

268

269 Date, K., Fagan, R., Crossland, S., Maceachern, D., Pyper, B., Bokanyi, R., Houze, Y., Andress, E., Tauxe, R.,  
 270 2011. Three outbreaks of foodborne botulism caused by unsafe home canning of vegetables-Ohio and  
 271 Washington, 2008 and 2009. J. Food Prot. 74(12), 2090–2096.

272

273 De Medici, D., Anniballi, F., Wyatt, G. M., Lindström, M., Messelhäusser, U., Aldus, C. F., Delibato, E.,  
 274 Korkeala, H., Peck, M. W., Fenicia, L., 2009. Multiplex PCR for detection of botulinum neurotoxin-producing  
 275 clostridia in clinical, food, and environmental samples. Appl Environ Microbiol. 75(20), 6457–6461.

276

- 277 Derman, Y., Lindström, M., Selby, K., Korkeala, H., 2011. Growth of group II *Clostridium botulinum* strains at  
278 extreme temperatures. J. Food Prot. 74(11), 1797–1804.  
279
- 280 ECFF (European Chilled Food Federation), 2006. Recommendations for the production of prepackaged  
281 chilled food. <https://www.ecff.net/best-practice/> (accessed 9 December 2019)  
282
- 283 Fenicia, L., Anniballi, F., Poushaban M., Franciosa, G., Aureli, P., 2002. Presence of *Clostridium botulinum*  
284 spores in sea-salt in Italy. Poster presented at the 18th International ICFHM Symposium, Food  
285 Microbiology, Lillehammer, Norway, 18–23. August 2002.  
286
- 287 Fernández, P.S., Peck, M.W., 1999. A predictive model that describes the effect of prolonged heating at 70  
288 to 90°C and subsequent incubation at refrigeration temperatures on growth from spores and toxigenesis by  
289 nonproteolytic *Clostridium botulinum* in the presence of lysozyme. Appl. Environ. Microbiol. 65(8), 3449–  
290 3457.  
291
- 292 Galazka, A., Przybylska, A., 1999. Surveillance of foodborne botulism in Poland: 1960–1998. Euro Surveill.  
293 4(6), 69–72.  
294
- 295 Gimenez, D.F., Gimenez, J.A., 1993. Serological subtypes of botulinal neurotoxins, in: DasGupta, B.R. (Eds.),  
296 Botulinum and tetanus neurotoxins. Plenum Press, New York, pp. 421–431.  
297
- 298 Gorris, L.G.M., Peck, M.V., 1998. Microbiological safety considerations when using hurdle technology with  
299 refrigerated processed foods of extended durability, in: Ghazala, S. (Eds.), sous vide and cook-chill  
300 processing for the food industry. Aspen publishers, Inc., Gaithersburg, pp. 207–233.  
301

- 302 Graham, A.F., Mason, D.R., Maxwell, F.J., Peck, M.W., 1997. Effect of pH and NaCl on growth from spores of  
 303 non-proteolytic *Clostridium botulinum* at chill temperature. Lett. Appl. Microbiol. 24(2), 95–100.  
 304
- 305 Hauschild H.W., Aris, B.J., Hilsheimer R, 1975. *Clostridium botulinum* in Marinated Products. Can. Inst. Food  
 306 Sci. Technol. J. 8(2), 84–87.  
 307
- 308 Hauschild, A.H., Hilsheimer, R., 1977. Enumeration of *Clostridium botulinum* spores in meats by a pour-  
 309 plate procedure. Can. J. Microbiol. 23(6), 829–832.  
 310
- 311 Hauschild, A.H.W., Hilsheimer, R., 1979. Effect of salt content and pH on toxigenesis by  
 312 *Clostridium botulinum* in Caviar. J. Food Prot. 42(3), 245–248.  
 313
- 314 Havlik, J., Plachy, V., Fernandez, J., Rada, V., 2010. Dietary purines in vegetarian meat analogues. J. Sci. Food  
 315 Agric. 90(14):2352 –2357.  
 316
- 317 Hellmich, D., Wartenberg, K.E., Zierz, S., Mueller, T.J., 2018. Foodborne botulism due to ingestion of home-  
 318 canned green beans: two case reports. J. Med. Case Rep. 12(1), 1.  
 319
- 320 Hielm, S., Hyytiä, E., Ridell, J., Korkeala, H., 1996. Detection of *Clostridium botulinum* in fish and  
 321 environmental samples using polymerase chain reaction. Int. J. Food Microbiol. 31(1–3), 357–365.  
 322
- 323 Hielm, S., Hyytiä, E., Andersin, A.B., Korkeala, H., 1998. A high prevalence of *Clostridium botulinum* type E in  
 324 Finnish freshwater and Baltic Sea sediment samples. J Appl Microbiol. 84(1):133 –137.  
 325

- 326 Hill, K.K., Smith, T.J., Helma, C.H., Ticknor, L.O., Foley, B.T., Svensson, R.T., Brown, J.L., Johnson, E.A., Smith,  
327 L.A., Okinaka, R.T., Jackson, P.J., Marks, J.D., 2007. Genetic diversity among botulinum neurotoxin-  
328 producing clostridial strains. J. Bacteriol. 189(3), 818–832.
- 329
- 330 Hill, S.E., Iqbal, R., Cadiz, C.L., Le, J., 2013. Foodborne botulism treated with heptavalent botulism antitoxin.  
331 Ann. Pharmacother. 47(2), e12.
- 332
- 333 Hyytiä, E., Hielm, S., Korkeala, H., 1998. Prevalence of *Clostridium botulinum* type E in Finnish fish and  
334 fishery products. Epidemiol Infect. 120(3), 245–250.
- 335
- 336 Insalata, N.F., Witzeman, S.J., Fredericks, G.J., Sunga, F.C.A., 1969. Incidence study of spores of  
337 *Clostridium botulinum* in convenience foods. Appl. Microbiol. 17(4), 542–544.
- 338
- 339 Jalava, K., Selby, K., Pihlajasaari, A., Kolho, E., Dahlsten, E., Forss, N., Bäcklund, T., Korkeala, H., Honkanen-  
340 Buzalski, T., Hulkko, T., Derman, Y., Järvinen, A., Kotilainen, H., Kultanen, L., Ruutu, P., Lyytikäinen, O.,  
341
- 342 Kalaycıoğlu, Z., Erım, F.B., 2019. Nitrate and nitrites in foods: worldwide regional distribution in view of  
343 their risks and benefits. J. Agric. Food Chem. 67(26), 7205–7222.
- 344
- 345 Kautter, D. A., 1964. *Clostridium botulinum* type E in smoked fish. J Food Sci. 29(6), 843–849.
- 346
- 347 Keto-Timonen, R., Heikinheimo, A., Eerola, E., Korkeala, H., 2006. Identification of *Clostridium species* and  
348 DNA fingerprinting of *Clostridium perfringens* by amplified fragment length polymorphism analysis. J Clin  
349 Microbiol. 44(11), 4057–4065.
- 350



- 351 Keto-Timonen, R., Lindström, M., Puolanne, E., Niemistö, M., Korkeala, H., 2012. Inhibition of toxigenesis of  
 352 group II (nonproteolytic) *Clostridium botulinum* type B in meat products by using a reduced level of  
 353 nitrite. *J. Food Prot.* 75(7), 1346–1349.  
 354
- 355 Kim, J., Foegeding, P.M., 1992. Principles of control, in: Hauschild, A.H.W., Dodds, K.L. (Eds.),  
 356 *Clostridium botulinum*: ecology and control in foods. Marcel Dekker Inc., New York, pp 121–176.  
 357
- 358 King, L.A., Niskanen, T., Junnikkala, M., Moilanen, E., Lindström, M., Korkeala, H., Korhonen, T., Popoff, M.,  
 359 Mazuet, C., Callon, H., Pihier, N., Peloux, F., Ichai, C., Quintard, H., Dellamonica, P., Cua, E., Lasfargue, M.,  
 360 Pierre, F., de Valk, H., 2009. Botulism and hot-smoked whitefish: a family cluster of type E botulism in  
 361 France, September 2009. *Euro Surveill.* 14(45), pii=19394.  
 362
- 363 Korkeala, H., Stengel, G., Hyytiä, E., Vogelsang, B., Bohl, A., Wihlman, H., Pakkala, P., Hielm, S., 1998. Type E  
 364 botulism associated with vacuum-packaged hot-smoked whitefish. *Int. J. Food Microbiol.* 43(1–2), 1–5.  
 365
- 366 Lilly, T.J., Solomon, H.M., Rhodehamel, E.J., 1995. Incidence of *Clostridium botulinum* in vegetables  
 367 packaged under vacuum or modified atmosphere. *J. Food Prot.* 59(1), 59–61.  
 368
- 369 Lindström, M., Keto, R., Markkula, A., Nevas, M., Hielm, S., Korkeala, H., 2001. Multiplex PCR assay for  
 370 detection and identification of *Clostridium botulinum* types A, B, E, and F in food and fecal material. *Appl.*  
 371 *Environ Microbiol.* 67(12), 5694–5699.  
 372
- 373 Lindström, M., Kiviniemi, K., Korkeala, H., 2006a. Hazard and control of group II (non-proteolytic)  
 374 *Clostridium botulinum* in modern food processing. *Int. J. Food Microbiol.* 108(1), 92–104.  
 375

- 376 Lindström, M., Vuorela, M., Hinderink, K., Korkeala, H., Dahlsten, E., Raahenmaa, M., 2006b. Botulism  
 377 associated with vacuum-packed smoked whitefish in Finland, June-July 2006. *Euro Surveill.* 11(29),  
 378 pii=3004.
- 379
- 380 Lindström, M., 2011. Two cases of food-borne botulism in Finland caused by conserved olives, October  
 381 2011. *Euro Surveill.* 16(49), 20034.
- 382
- 383 Lund, B.M., George, S. M., Franklin, J. G., 1987. Inhibition of type A and type B (proteolytic)  
 384 *Clostridium botulinum* by sorbic acid. *Appl Environ Microbiol.* 53(5), 935–941.
- 385
- 386 Lund, B.M., Peck, M.W., 1994. Heat resistance and recovery of spores of non-proteolytic *Clostridium*  
 387 *botulinum* in relation to refrigerated, processed foods with an extended shelf-life. *Journal of Applied*  
 388 *Bacteriology Symposium Supplement.* 76, 115–128.
- 389
- 390 MacDonald, K.L., Spengler, R.F., Hatheway, C.L., Hargrett, N.T., Cohen, M.L., 1985. Type A botulism from  
 391 sauteed onions. Clinical and epidemiologic observations. JAMA. 253(9), 1275–1278.
- 392
- 393 Mazuet, C., Silva, J.-D.N., Legeay, C., Sautereau, J., Popoff, R.M., 2018. Le botulisme humain en France,  
 394 2013–2016. Bull. Epidémiol. Hebd. 3, 46–54.
- 395
- 396 Meng, J., Genigeorgis, C.A., 1993. Modeling lag phase of nonproteolytic *Clostridium botulinum* toxigenesis  
 397 in cooked turkey and chicken breast as affected by temperature, sodium lactate, sodium chloride and spore  
 398 inoculum. *Int. J. Food Microbiol.* 19(2), 109–122.
- 399
- 400 Meng, J., Genigeorgis C.A. 1994. Delaying toxigenesis of *C. botulinum* by sodium lactate in 'sous-vide'  
 401 products. *Lett. Appl. Microbiol.* 19, 20–23.

402

403 Miller, A.J., Call, J.E., Whiting, R.C., 1993. Comparison of organic acid salts for *Clostridium botulinum* control  
404 in an uncured turkey product. J. Food Prot. 56(11), 958–962.

405

406 Morse, D.L., Pickard, L.K., Guzewich, J.J., Devine, B.D., Shayegani, M., 1990. Garlic-in-oil associated  
407 botulism: episode leads to product modification. Am. J. Public Health. 80(11), 1372–1373.

408

409 Peck, M.W., Fairbairn, D.A., Lund, B.M., 1993. Heat-resistance of spores of non-proteolytic  
410 *Clostridium botulinum* estimated on medium containing lysozyme. Lett. Appl. Microbiol. 16(3), 126–131.

411

412 Peck, M.W., Fernandez, P.S., 1995. Effect of lysozyme concentration, heating at 90 degrees °C, and then  
413 incubation at chilled temperatures on growth from spores of non-proteolytic *Clostridium botulinum*. Lett  
414 Appl. Microbiol. 21(1), 50–54.

415

416 Peck, M.W., Lund, B.M., Fairbairn, D.A., Kaspersson, A.S., Undeland, P.C., 1995. Effect of heat treatment on  
417 survival of, and growth from, spores of nonproteolytic *Clostridium botulinum* at refrigeration temperatures.  
418 Appl. Environ. Microbiol. 61(5), 1780–1785.

419


420 Peck, M.W., 2006. *Clostridium botulinum* and the safety of minimally heated, chilled foods: an emerging  
421 issue? J. Appl. Microbiol. 101(3), 556–570.

422

423 Segner, W.P., Schmidt, C.F., Boltz, J.K., 1966. Effect of sodium chloride and pH on the outgrowth of spores  
424 of type E *Clostridium botulinum* at optimal and suboptimal temperatures. Appl. Microbiol. 14(1), 49–54.

425

- 426 Sevenier, V., Delannoy, S., André, S., Fach, P., Remize, F., 2012. Prevalence of *Clostridium botulinum* and  
 427 thermophilic heat-resistant spores in raw carrots and green beans used in French canning industry. Int. J.  
 428 Food Microbiol. 155(3), 263–268.  
 429
- 430 Seward, R.A., Deibel, R.H., Lindsay, R.C., 1982. Effects of potassium sorbate and other antibotulinal agents  
 431 on germination and outgrowth of *Clostridium botulinum* type E spores in microcultures. Appl Environ  
 432 Microbiol. 44(5), 1212–1221.  
 433
- 434 Sobel, J., Tucker, N., Sulka, A., McLaughlin, J., Maslanka, S., 2004. Foodborne botulism in the United States,  
 435 1990–2000. Emerg. Infect. Dis. 10(9), 1606–1611.  
 436
- 437 Thomas, H.A., 1942. Bacterial densities from fermentation tube tests. J. Am. Water Works Assoc. 34(4),  
 438 572–576.  
 439
- 440 Woodburn, M., Somers, E., Rodriguez, J., Schantz, E., 1979. Heat inactivation rates of botulinum toxins A B E  
 441 and F in some foods and buffers. J. Food Sci. 44(6): 1658–1661.  
 442
- 443 Young-Perkins, K.E., Merson, L., 1987. *Clostridium botulinum* spore germination, outgrowth, and toxin  
 444 production below pH 4.6; interactions between pH, total acidity, and buffering capacity. J. Food Sci 52(4),  
 445 1084–1088.  
 446
- 447 Zanon, P., Pattis, P., Pittscheider, W., Roscia, G., De Giorgi, G., Sacco, G., Vötter, K., Stockner, I., De Giorgi,  
 448 F., Wiedermann, C.J., 2006. Two cases of foodborne botulism with home-preserved asparagus. Anesthesiol.  
 449 Intensivmed. Notfallmed. Schmerzther. 41(3), 156–159.  
 450  
 451



## 459 FIGURE LEGEND

460 Fig.1. Dendrogram of eight *Clostridium botulinum* isolates originating from vegetarian sausages and 25 *C.*  
461 *botulinum* strains included in Keto-Timonen et al. (2006) based on AFLP analysis. A similarity analysis was  
462 performed using the Pearson product-moment correlation coefficient, and clustering was performed by  
463 using the unweighted pair-group method with arithmetic averages. Isolates originating from vegetarian  
464 sausages are written in bold font. <sup>a</sup>Previously published by Keto-Timonen et al. (2006).

465 Table 1. The prevalence of *Clostridium botulinum* and BoNT genes in vegetarian sausages.


Product type	No. of samples examined	No. of positive samples (%)	MPN estimate of <i>C. botulinum</i> cell count (cells/kg)	No. of samples positive for one or two BoNT genes (% of positive samples)					
				Type A	Type B	Type E	Type F	Types B and E <sup>a</sup>	Types B and F <sup>a</sup>
Vacuum-packaged	66	23 (35%)	20–1200	8 (35%)	7 (30%)	6 (26%)	ND	1 (4%)	1 (4%)
Frozen	8	1 (13%)	110	ND	1 (100%)	ND	ND	ND	ND
Total	74	24 (32%)	20–1200	8 (33%)	8 (33%)	6 (25%)	ND	1 (4%)	1 (4%)

466 <sup>a</sup>Both types detected in the same sample.

467 ND, not detected.

Journal Pre-proof





- High prevalence of *Clostridium botulinum* found in vegetarian sausage products
- *C. botulinum* Groups I and II, and genes for neurotoxins A, B, E, and F were found
- *C. botulinum* Group II is the main food safety concern in chilled packaged foods

Declarations of interest: none.

Journal Pre-proof